

adenosylcobyric acid synthase (glutamine-hydrolysing)

Cat. No. EXWM-5803

Lot. No. (See product label)

Introduction

Description Requires Mg2+. NH3 can act instead of glutamine. This enzyme catalyses the four-step amidation

sequence from cobyrinic acid a,c-diamide to cobyric acid via the formation of cobyrinic acid triamide,

tetraamide and pentaamide intermediates.

Synonyms CobQ; cobyric acid synthase; 5'-deoxy-5'-adenosylcobyrinic-acid-a,c-diamide:L-glutamine amido-ligase;

Ado-cobyric acid synthase [glutamine hydrolyzing]

Product Information

Form Liquid or lyophilized powder

EC Number EC 6.3.5.10

CAS No. 137672-90-3

Reaction 4 ATP + adenosylcobyrinic acid a,c-diamide + 4 L-glutamine + 4 H2O = 4 ADP + 4 phosphate +

adenosylcobyric acid + 4 L-glutamate

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce

according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C \sim -80 °C.

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1