

Transglutaminase 2 from Human tissue, Recombinant

Cat. No. NATE-1728

Lot. No. (See product label)

Introduction

Description	This enzyme belongs to the family of transferases, specifically those transferring phosphorus-containing groups (phosphotransferases) with a phosphate group as acceptor.
Applications	Transglutaminase 2 catalyzes acyl transfer reactions from glutamin residues in proteins or peptides to primary amines, e. g. the formation of ϵ -(γ -glutamyl) lysine bonds between proteins by transferring the acyl group of a peptide-bound glutamine residue to the primary amino group of a peptide-bound lysine residue. Transglutaminase 2 may also be used for immunoprecipitation.
Synonyms	transglutaminase; EC 2.3.2.13; 80146-85-6; transglutaminase; Factor XIIIa; fibrinolygase; fibrin stabilizing factor; glutaminylpeptide γ -glutamyltransferase; polyamine transglutaminase; tissue transglutaminase; R-glutaminyl-peptide:amine γ -glutamyl transferase; protein-glutamine γ -glutamyltransferase; TG1

Product Information

Species	Human
Source	Insect cells
Appearance	White lyophilized solid.
Form	The transglutaminase is lyophilized from 10 mM Tris-HCl pH 8.1, 150 mM NaCl, 1 mM EDTA, 5 mM DTT. Sample contains maltodextrin.
EC Number	EC 2.3.2.13
CAS No.	80146-85-6
Molecular Weight	78 kDa
Purity	> 90 % (visually by SDS-PAGE)
Activity	> 1500 U/mg [Activity is determined by measuring the rate of fluorescence enhancement after transglutaminase-catalyzed monodansylcadaverine-incorporation into N,N-dimethylated casein according to Lorand et al., Anal. Biochem. 44 (221-231).
Activators	Add 10 mM Ca ²⁺ to activate transglutaminase.
Unit Definition	1 U is defined as the increase in fluorescence intensity of 1 a.u./min (measured on a Cary eclipse fluorescence spectrophotometer, Varian; λ_{ex} = 332 nm, λ_{em} = 500 nm; band filter = 5 nm; detector strength = 600 V; temperature = 37°C, assay volume = 1 ml)].

Usage and Packaging

Package	250 μ g; 1mg
Reconstitution	Add the volume of water specified in the certificate of analysis under aliquotation to the vial of lyophilized powder. Rotate vial gently until solid dissolves. After reconstitution the solution should be stored frozen in working aliquots.

Storage and Shipping Information

Storage Store working aliquots at $\leq -20^{\circ}\text{C}$. Avoid repeated freezing and thawing. Delivery is possible at ambient temperature.