

## Transglutaminase 2 from Dog, Recombinant

Cat. No. NATE-1733

Lot. No. (See product label)

### Introduction

<b>Description</b>	This enzyme belongs to the family of transferases, specifically those transferring phosphorus-containing groups (phosphotransferases) with a phosphate group as acceptor.
<b>Applications</b>	The transglutaminase 2 catalyzes acyl transfer reactions from glutamine residues in proteins or peptides to primary amines, e. g. the formation of $\epsilon$ -( $\gamma$ -glutamyl) lysine bonds between proteins by transferring the acyl group of a peptide-bound glutamine residue to the primary amino group of a peptide-bound lysine residue. The transglutaminase 2 may also be used for immunoprecipitation.
<b>Synonyms</b>	transglutaminase; EC 2.3.2.13; 80146-85-6; transglutaminase; Factor XIIIa; fibrinolygase; fibrin stabilizing factor; glutaminylpeptide $\gamma$ -glutamyltransferase; polyamine transglutaminase; tissue transglutaminase; R-glutaminyl-peptide:amine $\gamma$ -glutamyl transferase; protein-glutamine $\gamma$ -glutamyltransferase; TG1

### Product Information

<b>Species</b>	Dog
<b>Source</b>	Insect cells
<b>Appearance</b>	White lyophilized solid.
<b>Form</b>	The transglutaminase is lyophilized from 10 mM Tris-HCl pH 8.1, 150 mM NaCl, 1 mM EDTA, 5 mM DTT. Sample contains maltodextrin.
<b>EC Number</b>	EC 2.3.2.13
<b>CAS No.</b>	80146-85-6
<b>Molecular Weight</b>	78 kDa
<b>Purity</b>	> 90 % (visually by SDS-PAGE)
<b>Activity</b>	> 1500 U/mg [Activity is determined by measuring the rate of fluorescence enhancement after transglutaminase-catalyzed monodansylcadaverine-incorporation into N,N-dimethylated casein according to Lorand et al., Anal. Biochem. 44 (221-231).
<b>Activators</b>	Add 10 mM Ca <sup>2+</sup> to activate transglutaminase.
<b>Unit Definition</b>	1 U is defined as the increase in fluorescence intensity of 1 a.u./min (measured on a Cary eclipse fluorescence spectrophotometer, Varian; $\lambda_{ex}$ = 332 nm, $\lambda_{em}$ = 500 nm; band filter = 5 nm; detector strength = 600 V; temperature = 37°C, assay volume = 1 ml)].

### Usage and Packaging

<b>Package</b>	250 $\mu$ g; 1mg
<b>Reconstitution</b>	Add the volume of H <sub>2</sub> O the protein is lyophilized from (see Certificate of Analysis) to the vial of lyophilized powder. Rotate vial gently until solid dissolves. After reconstitution the solution should be stored frozen in working aliquots. Keep cooled on ice for short term storage.

### ***Storage and Shipping Information***

**Storage** Store working aliquots at  $\leq -20^{\circ}\text{C}$ . Avoid repeated freezing and thawing. Delivery is possible at ambient temperature.