

## **Native Bacillus sp Chitosanase**

Cat. No. NATE-1746 Lot. No. (See product label)

Introduction	
Description	Chitosanase is a powdered chitosanase preparation made by submerged fermentation of a selected strain of the bacterium Bacillus sp. The enzyme catalyzes the breakdown of chitosan, a partially or completely de-acetylated derivative of chitin ( $\beta$ -1,4 homopolymer of N-acetyl glucosamine).
Applications	Chitosanase can be used for hydrolyzing chitosan(degree of de-acetylatin: $40 \sim 100\%$ ). Especially, it can be used for the production of chitosan oligosaccharides from chitosan, which have a variety of biological activities such as immunostimulating activity, anti-tumor activity, anti-microbial activity, etc.
Synonyms	Chitosanase; EC 3.2.1.132; 51570-20-8; Chitosan N-acetylglucosaminohydrolase
Product Information	
Source	Bacillus sp
Appearance	White or light yellow colored, freeze-dried powder
EC Number	EC 3.2.1.132
CAS No.	51570-20-8
Molecular Weight	45,000Da estimated by SDS-PAGE
Activity	35,000U/g
pH Stability	Stable in pH range of 4.5 to 8.0
Optimum pH	pH range of 4.5 to 6.0
Thermal stability	More than 90% activity remains after 24 hr incubation at 40°C.
Optimum temperature	60°C
Specificity	Shows high activities against chitosan substrates which are de-acetylated by 40 to 100%.
Unit Definition	The standard activity is determined by modified Schales method. One unit(U) is defined as the amount of enzyme that releases one µmole of reducing sugar (measured as D-glucosamine equivalents) from chitosan per minute at pH 5.0 at 48°C. A detailed description of the method is available on request.
Notes	Arsenic: Less than 4ppm. Heavy metals: Less than 10ppm (as Pb). Coli-form bacteria: Less than 30 colony-forming units(CFU) per gram. Samonella: Negative.

## Storage and Shipping Information

Storage

The product should be stored in a cool, dry environment with temperatures below 4°C.