

N-Acetylglucosaminyltransferases (Crude Enzyme)

Cat. No. NATE-1813 Lot. No. (See product label)

Introduction	
Description	This product with the indicated enzyme activity was briefly purified from engineered E. coli.
Applications	biotechnology; medicine; pharmacology
Synonyms	N-acetylglucosaminyltransferase V; α -mannoside β -1,6-N- acetylglucosaminyltransferase; uridine diphosphoacetylglucosamine- α -mannoside β 1 \rightarrow 6-acetylglucosaminyltransferase; UDP-N-acetylglucosamine: α -mannoside- β 1,6 N-acetylglucosaminyltransferase; α -1,3(6)-mannosylglycoprotein β -1,6-N- acetylglucosaminyltransferase; GnTV
Product Information	
Source	E. coli
Appearance	Clear to translucent yellow solution
EC Number	EC 2.4.1.155
CAS No.	83588-90-3
Activity	Undetermined
Reaction	UDP-N-acetyl-D-glucosamine + 6-(2-[N-acetyl-β-D-glucosaminyl]-α-D-mannosyl)-β- D-mannosyl-R = UDP + 6-(2,6-bis[N-acetyl-β-D-glucosaminyl]-α-D-mannosyl)-β-D- mannosyl-R
Notes	Since this product needs to be freshly prepared, it will take about 2 weeks after you confirm the order. Each time of the freeze-thawing may cause partial inactivation. Therefore, it should be dispensed as required and stored at -20 ° C or lower. With the preservation of the extension of time, the enzyme activity will decline to a certain extent, so the product should be used as soon as possible. This product may have turbidity or precipitation in the production and preservation process, it can be mixed after melting and will not affect the normal use. This product is limited to scientific research use, shall not be used for clinical diagnosis or treatment, shall not be used for food or medicine, shall not be stored in ordinary residential. For your safety and health, please wear an experimental suit and wear disposable gloves.
Usage and Packaging	
Package	100ml
Storage and Shipping Information	
Storage	at -20 °C or lower, for at least 1 month.