

## Nicotinamide Mononucleotide Adenylyltransferase (Crude Enzyme)

Cat. No. NATE-1830

Lot. No. (See product label)

### Introduction

**Description** This enzyme belongs to the family of transferases, specifically those transferring phosphorus-containing nucleotide groups (nucleotidyltransferases). This enzyme participates in nicotinate and nicotinamide metabolism. The human version of this protein is NMNAT1. This product with the indicated enzyme activity was briefly purified from engineered E. coli.

**Applications** drug development; medicine

**Synonyms** NAD + pyrophosphorylase; adenosine triphosphate-nicotinamide mononucleotide transadenylase; ATP:NMN adenylyltransferase; diphosphopyridine nucleotide pyrophosphorylase; nicotinamide adenine dinucleotide pyrophosphorylase; nicotinamide mononucleotide adenylyltransferase; NMN adenylyltransferase

### Product Information

**Source** E. coli

**Appearance** Clear to translucent yellow solution

**EC Number** EC 2.7.7.1

**CAS No.** 9032-70-6

**Activity** Undetermined

**Reaction** ATP + nicotinamide ribonucleotide = diphosphate + NAD +

**Notes** Since this product needs to be freshly prepared, it will take about 2 weeks after you confirm the order. Each time of the freeze-thawing may cause partial inactivation. Therefore, it should be dispensed as required and stored at -20 °C or lower. With the preservation of the extension of time, the enzyme activity will decline to a certain extent, so the product should be used as soon as possible. This product may have turbidity or precipitation in the production and preservation process, it can be mixed after melting and will not affect the normal use. This product is limited to scientific research use, shall not be used for clinical diagnosis or treatment, shall not be used for food or medicine, shall not be stored in ordinary residential. For your safety and health, please wear an experimental suit and wear disposable gloves.

### Usage and Packaging

**Package** 100ml

### Storage and Shipping Information

**Storage** at -20 °C or lower, for at least 1 month.