

Native *Bacillus stearothermophilus* Diaphorase 1

Cat. No. NATE-1901

Lot. No. (See product label)

Introduction

Description

Diaphorase catalyzes the reaction of a reduced di- or tri-phosphopyridine nucleotide hydrogen donor with a hydrogen acceptor, usually a dye in the leucoform.

Applications

The enzyme is useful for the measurement of various dehydrogenase reactions in visible spectral range.

Synonyms

Diaphorase 1; Di-1; EC 1.6.99 -

Product Information

Source

Bacillus stearothermophilus

Appearance

Lyophilized

EC Number

EC 1.6.99.-

Molecular Weight

ca. 30,000

Specific Activity

more than 1,000 U/mg protein

Contaminants

(as Diaphorase activity = 100 %) Adenylate kinase: < 0.01 %; NADH oxidase: < 0.01 %.

pH Stability

7.5 - 9.5

Optimum pH

8

Thermal stability

No detectable decrease in activity up to 50 °C.

Unit Definition

One unit of activity is defined as the amount of Di-1 that reduces 1 µmol of DCIP per minute at 30 °C.

Reaction

$\text{NAD(P)H} + \text{Acceptor(ox.)} + \text{H}^+ \leftrightarrow \text{NAD(P)}^+ + \text{Acceptor(red.)}$

Storage and Shipping Information

Storage

Stable at -20 to 5 °C for at least one year.