

## Protein Kinase Cı, Active human, Recombinant

Cat. No. NATE-0577

Lot. No. (See product label)

## Introduction

Description

Protein Kinase C (PKC) is a serine/threonine kinase that is activated intracellularly by signal transduction pathways that produce DAG from phosphatidylinositol diphosphate (PIP2) and phosphatidylcholine (PC) through the action of various activated phospholipases. Phorbol esters also stimulate PKC. At least 11 PKC isozymes have been identified that differ in primary structure, tissue distribution, subcellular localization, response to extracellular signals, and substrate specificity. The isozymes can be grouped into three subfamilies. Members of the first family require Ca2+ and phospholipid and include PKC $\alpha$ ,  $\beta$ I,  $\beta$ II, and  $\gamma$ . Members of the second family are phospholipid-dependent but Ca2+-independent, and include PKC $\alpha$ ,  $\beta$ I, and  $\beta$ I. Members of the third family are not activated by either DAG or phorbol esters and include PKC $\alpha$ ,  $\beta$ I, and  $\beta$ I.

**Applications** 

Kinase activity is measured as the molar amount of phosphate incorporated into the CREBtide substrate peptide per minute per mg protein at  $30^{\circ}$ C using a final concentration of  $50 \mu$ M [32P] ATP.

Synonyms

PKCL; Protein Kinase C Lambda/Iota; PKCı

## **Product Information**

**Species** Human

**Source** E. coli

**Form** buffered aqueous glycerol solution

Molecular

Weight

apparent mol wt ~98 kDa

**Purity** > 85% (SDS-PAGE)

Buffer Solution of 5 μg in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.25 mM DTT, 0.1 mM EGTA, 0.1mM EDTA, 0.1

mM PMSF, and 25% glycerol.

Pathway Cell junction organization, organism-specific biosystem; Cell-Cell communication, organism-specific

biosystem; Cell-cell junction organization, organism-specific biosystem; EGFR1 Signaling Pathway, organism-specific biosystem; Endocytosis, organism-specific biosystem; Endocytosis, conserved

biosystem; G Protein Signaling Pathways, organism-specific biosystem

Function ATP binding; metal ion binding; nucleotide binding; phospholipid binding; protein binding; protein kinase

C activity; protein kinase activity; protein serine/threonine kinase activity; protein serine/threonine

kinase activity; zinc ion binding

## Storage and Shipping Information

*Stability* −70°C

**Tel:** 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1