

## **ZAP-70 Tyrosine Kinase human, Recombinant**

Cat. No. NATE-0738

Lot. No. (See product label)

## Introduction

**Description** ZAP-70 is protein tyrosine kinase of the Syk family that is localized exclusively to

the cytosol of T cells and natural killer cells. It is required for T cell activation. It is activated by Lck-mediated phosphorylation of its tyrosine residues. ZAP-70 Tyrosine Kinase is a protein belongs to Syk family and is essential for T cell activation. It facilitates the upregulation of Fas ligand in activation-induced T cell apoptosis. Additionally, it also plays a pivotal role in LFA-1-dependent T Cell

Migration.

**Applications** Human ZAP-70 Tyrosine Kinase was used for mapping the phosphorylation sites on

LAT (linker for activation of T cells) for investigating the mechanism of activation of

signalling proteins in T cells.

**Synonyms** ZAP-70 Tyrosine Kinase; Zeta-chain-associated protein kinase 70; ZAP-70; SRK;

STCD; STD; TZK; ZAP70

## **Product Information**

**Species** Human

**Source** baculovirus infected insect cells (Histidine tagged)

**Form** buffered aqueous glycerol solution

**Molecular Weight** mol wt 70 kDa

*Purity* > 90% (SDS-PAGE)

**Activity** > 150 U/mg

**Buffer** Solution in 25 mM HEPES, pH 7.5, containing 150 mM NaCl, 0.5 mM DTT, and 50%

glycerol.

**Pathway** Adaptive Immune System, organism-specific biosystem; B Cell Receptor Signaling

Pathway, organism-specific biosystem; Generation of second messenger molecules, organism-specific biosystem; Immune System, organism-specific biosystem; Inflammatory Response Pathway, organism-specific biosystem; Natural killer cell mediated cytotoxicity, organism-specific biosystem; Natural killer cell mediated

cytotoxicity, conserved biosystem

**Function** ATP binding; non-membrane spanning protein tyrosine kinase activity; nucleotide

binding; protein binding; protein tyrosine kinase activity; protein tyrosine kinase

activity; protein tyrosine kinase activity

Unit Definition One unit will phosphorylate 1 nmol polyglutamic acid:tyrosine per min at 37°C at

pH 7.0.

## Storage and Shipping Information

*Stability* –70°C

**Tel:** 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com 1/1