

## Native *Bacillus stearothermophilus* Phosphofructokinase

Cat. No. NATE-0551

Lot. No. (See product label)

### Introduction

#### Description

Phosphofructokinase is a kinase enzyme that phosphorylates fructose 6-phosphate in glycolysis. The enzyme-catalysed transfer of a phosphoryl group from ATP is an important reaction in a wide variety of biological processes. One enzyme that utilizes this reaction is phosphofructokinase (PFK), which catalyses the phosphorylation of fructose-6-phosphate to fructose-1,6-bisphosphate, a key regulatory step in the glycolytic pathway. PFK exists as a homotetramer in bacteria and mammals (where each monomer possesses 2 similar domains) and as an octomer in yeast (where there are 4 alpha-(PFK1) and 4 beta-chains (PFK2), the latter, like the mammalian monomers, possessing 2 similar domains[3]). This protein may use the morpheein model of allosteric regulation.

#### Applications

Useful for enzymatic determination of fructose-6-phosphate

#### Synonyms

PFKWII; EC 2.7.1.11; PFK; phosphofructokinase; 6-phosphofructokinase; Phosphofructokinase I; Phosphohexokinase

### Product Information

#### Source

*Bacillus stearothermophilus*

#### Appearance

White to pale yellow powder

#### Form

Freeze dried powder

#### EC Number

EC 2.7.1.11

#### CAS No.

9001-80-3

#### Molecular Weight

72 kDa (gel filtration); 35 kDa (SDS-PAGE)

#### Activity

> 250 U/mg

#### Contaminants

NADPH oxidase < 0.01%; ATPase < 0.005%

#### Isoelectric point

pH 5.9

#### pH Stability

6.0-10.0 (37°C, 60 mins)

#### Optimum pH

9

#### Thermal stability

Stable at 55°C and below (pH 8.5, 30 mins)

#### Michaelis Constant

D-Fructose-6-phosphate (D-F-6-P) 5.8 mM (at 37°C); ATP 0.07 mM (at 37°C)

#### Activators

Mg<sup>2+</sup>

#### Unit Definition

One unit is defined as the amount of enzyme which converts 1 μmole of fructose-6-phosphate to Fructose-1,6-bisphosphate per minute at 37°C under the conditions specified in the assay procedure.

### Storage and Shipping Information

#### Storage

Storage at -20°C in the presence of a desiccant is recommended

