

Lacto-N-Biosidase, Recombinant

Cat. No. NATE-0855

Lot. No. (See product label)

Introduction

Description	Lacto-N-Biosidase may be used during glycoprotein analysis to better study the structure and function of glycoprotein and glycolipid sugar chains. This product can be used for specific oligosaccharide hydrolysis of type-I chain oligosaccharides, producing lacto-N-biose (Gal β 1-3 GlcNAc) as a byproduct, but not oligosaccharide hydrolysis of type-II chain oligosaccharides. As a result, Lacto-N-Biosidase can be used to distinguish type-I versus type-II glycoprotein as well as glycolipid sugar chains. When used in conjunction with α -1,3/4-Fucosidase, Lacto-N-Biosidase can also help distinguish Sialyl-Lewis x and Sialyl-Lewis a structures.
Applications	Specific hydrolysis of oligosaccharides with type-I sugar chains and production of byproduct lacto-N-biose (Gal β 1-3 GlcNAc)
Synonyms	Oligosaccharide lacto-N-biosylhydrolase; Lacto-N-Biosidase

Product Information

Source	Streptomyces sp. 142
Form	Solution in 50 mM sodium acetate buffer, pH 5.5, containing 0.05% Brij-58
Molecular Weight	60 kDa (SDS-PAGE)
Concentration	1 μ U/ μ l
pH Stability	pH 4.0–10 (4°C, 16 hrs.)
Optimum pH	pH 5.5
Unit Definition	One unit is the amount of enzyme required to hydrolyze 1 μ mol of PA-lacto-N-tetraose in 1 minute at 37°C, pH 5.5.