

## Native Bovine Angiotensin Converting Enzyme

Cat. No. NATE-0866

Lot. No. (See product label)

### Introduction

#### Description

Angiotensin-converting enzyme (EC 3.4.15.1), or "ACE" indirectly increases blood pressure by causing blood vessels to constrict. It does that by converting angiotensin I to angiotensin II, which constricts the vessels. For this reason, drugs known as ACE inhibitors are used to lower blood pressure. ACE, angiotensin I and angiotensin II are part of the renin-angiotensin system (RAS), which controls blood pressure by regulating the volume of fluids in the body. ACE is secreted in the lungs and kidneys by cells in the endothelium (inner layer) of blood vessels.

#### Applications

Positive Control

#### Synonyms

ACE; Angiotensin Converting Enzyme; EC 3.4.15.1; dipeptidyl carboxypeptidase I; peptidase P; dipeptide hydrolase, peptidyl dipeptidase; angiotensin converting enzyme; kininase II; angiotensin I-converting enzyme; carboxycathepsin; dipeptidyl carboxypeptidase; "hypertensin converting enzyme" peptidyl dipeptidase I; peptidyl-dipeptide hydrolase; peptidyl dipeptide hydrolase; endothelial cell peptidyl dipeptidase; peptidyl dipeptidase-4; PDH; peptidyl dipeptide hydrolase; DCP

### Product Information

#### Species

Bovine

#### Source

Bovine lung

#### EC Number

EC 3.4.15.1

#### Purity

>95% as determined by SDS-PAGE.

#### Activity

One unit produces 1 mmole of hippuric acid or His-Leu from Hippuril-His-Leu per min in 0.1 M phosphate buffer-300 mM NaCl, pH 8.3 at 37°C.

#### Optimum pH

pH 7.5-8.5

### Storage and Shipping Information

#### Storage

Maintain at -20°C in undiluted aliquots for up to 12 months. Maintain at 4°C for up to one month. A decrease in activity may occur within prolonged storage at 4°C.