

Native Staphylococcus aureus Endoproteinase GluC

Cat. No. NATZ-137

Lot. No. (See product label)

Introduction

Description

This enzyme is isolated from Staphylococcus aureus V8 and specifically cleaves at the carboxyl side of glutamic acid residues. It cleaves peptide bonds at the carboxyl side of both glutamic acid and aspartic acid in phosphate buffer systems, and at the carboxyl side of glutamic acid in ammonium buffer systems. Due to its high substrate specificity, it is used for peptide mapping in various protein sequencing analyses. 1. Cleavage Exclusively at the Carboxyl Terminus of Glutamic Acid Digest at 30–37°C for 2–24 hours in either 0.01–0.1 mol/L ammonium bicarbonate (pH 7.8) or 0.01–0.1 mol/L ammonium acetate (pH 4.0), with an enzyme-to-substrate molar ratio of 1:30–100. 2. Simultaneous Cleavage at the Carboxyl Terminus of Aspartic Acid Digest under the same conditions as above in 0.01–0.1 mol/L phosphate buffer (pH 7.8).

Applications

Specifically cleaves at the carboxyl side of glutamic acid residues. Peptide mapping for sequence analysis of various proteins.

Synonyms

Protease V8

Product Information

Source

Staphylococcus aureus V8

Form

White crystals or crystalline powder

CAS No.

66676-43-5

Molecular Weight

Approximately 27,000

Activity

20 units/mg or higher

Optimum pH

Proteolytic activity is detectable across the pH range of 3.5–9.5, with maximum activity observed at pH 4.0 and pH 7.8 (using hemoglobin as the substrate).

Unit Definition

One unit is defined as the amount of enzyme that produces 1 µmol of 4-nitroaniline from the substrate 2-Phe-Leu-Glu-4-NA per minute at pH 7.8 and 25°C.