

## **Native Human Cancer Antigen 19-9**

Cat. No. NATE-0956

Lot. No. (See product label)

## Introduction

**Description** Cancer Antigen 19-9 is a tumor marker elevated in blood of patients with

carcinoma of the gastro-intestinal tract. Primarily used distinguishing pancreatic cancer from pancreatitis, CA 19-9 is not sufficiently specific for use as a cancer screening test. The specificity for pancreatic cancer increases with increasing levels, high levels showing a specificity of >97% for tumor presence. CA 19-9 can be elevated in many types of gastrointestinal cancer, such as colorectal cancer, esophageal cancer and hepatocellular carcinoma. A group of mucin type

esophageal cancer and hepatocellular carcinoma. A group of mucin type glycoprotein Sialosyl Lewis Antigens (SLA), such as CA19-9 and CA19-5, have come to be recognized as circulating cancer associated antigens for gastrointestinal cancer. CA19-9 represents the most important and basic carbohydrate tumor marker. The immunohistologic distribution of CA19-9 in tissues is consistent with the quantitative determination of higher CA19-9 concentrations in cancer than in normal or inflamed tissues. Recently reports indicates that the serum CA19-9 level is frequently elevated in the serum of subjects with various gastrointestinal malignancies, such as pancreatic, colorectal, gastric and hepatic carcinomas. Together with CEA, elevated CA19-9 is suggestive of gallbladder neoplasm in the

setting of inflammatory gallbladder disease.

Applications Diagnostic Controls; Calibrators & Standards; Immunoassays; Testing/Assay

Validation; Life Science; Validation Studies; Manufacturing; Tumor Markers

**Synonyms** Tumor Marker 19-9; CA 19-9; Cancer Antigen 19-9; carbohydrate antigen 19-9;

Sialylated Lewis (a) Antigen

## **Product Information**

**Species** Human

**Source** Human Liver Metastases

**Appearance** Clear and Colorless

Form Liquid in phosphate buffered saline, pH 7.4 with sucrose and 0.05% sodium azide

**Purity** High Purity - Low Cross Contamination of other cancer antigens (Gel filtration & ion-

exchange chromatography).

**Activity** Typically > 500 kU/mL

## Storage and Shipping Information

**Storage** Store at -20°C

**Stability** 2 years

1/1