

Glutamate Dehydrogenase (NAD(P)) from E.coli, Recombinant

Cat. No. NATE-0981 Lot. No. (See product label)

Introduction

Description	Glutamate dehydrogenase (GLDH) is an enzyme, present in most microbes and the mitochondria of eukaryotes, as are some of the other enzymes required for urea synthesis, that converts glutamate to α -ketoglutarate, and vice versa. In animals, the produced ammonia is usually used as a substrate in the urea cycle. Typically, the α -ketoglutarate to glutamate reaction does not occur in mammals, as glutamate dehydrogenase equilibrium favours the production of ammonia and α -ketoglutarate.
Applications	Use recombinant Glutamate Dehydrogenase in diagnostic tests for the determination of ammonia, urea, L-glutamate, glutamate pyruvate transaminase and leucine aminopeptidase.
Synonyms	glutamate dehydrogenase (NADP+); glutamic dehydrogenase; dehydrogenase; glutamate (nicotinamide adenine dinucleotide (phosphate)); glutamic acid dehydrogenase; L-glutamate dehydrogenase; L-glutamic acid dehydrogenase; NAD(P)-glutamate dehydrogenase; NAD(P)H-

dependent glutamate dehydrogenase; glutamate dehydrogenase (NADP); GLDH

Product Information

Source	E.coli
Appearance	White lyophilizate
CAS No.	2604152
Molecular Weight	\sim 2 200 kD for the associated enzyme with 8 subunits; 280 kD for one subunit.
Activity	>80 U/mg
Contaminants	Alcohol dehydrogenase: <0.005 Lactate dehydrogenase: <0.005 Malate dehydrogenase: <0.005 "NADH-Oxidase": <0.005 NH4: <0.05 μg/mg lyophilizate
pH Stability	5.5-6.5
Optimum pH	8
Michaelis	I-glutamate: 1.8 x 10-3 mol/l NADP: 4.7 x 10-5 mol/l α -ketoglutarate: 7.0 x 10-4 mol/l NH4+: 3.2 x 10-3
Constant	mol/l NADPH: 2.6 x 10-5 mol/l Km values for NAD or NADH are difficult to obtain due to their inhibitory action.
<i>Constant</i>	mol/l NADPH: 2.6 x 10-5 mol/l Km values for NAD or NADH are difficult to obtain due to their inhibitory action. The oxidation of L-glutamate is stimulated by ADP and inhibited by GTP. In contrast, the oxidation of alanine, leucine, isoleucine, methionine, valine, norleucine, norvaline and 2-aminobutyrates is stimulated by ADP.
<i>Constant</i> <i>Specificity</i> <i>Activators</i>	 mol/l NADPH: 2.6 x 10-5 mol/l Km values for NAD or NADH are difficult to obtain due to their inhibitory action. The oxidation of L-glutamate is stimulated by ADP and inhibited by GTP. In contrast, the oxidation of alanine, leucine, isoleucine, methionine, valine, norleucine, norvaline and 2-aminobutyrates is stimulated by GTP and inhibited by ADP. Thioglycolic acid, b-mercaptoethylamine, EDTA, α, α'-dipyridyl

Storage and Shipping Information

Stability At +2 to +8°C within specification range for 12 months. Store dry.