

## Puromycin Dihydrochloride

Cat. No. CEI-0634

Lot. No. (See product label)

### Introduction

#### Description

A nucleoside antibiotic. It inhibits protein synthesis by disrupting peptide transfer on ribosomes causing premature chain termination during translation. It is a potent translational inhibitor in both prokaryotic and eukaryotic cells. Resistance to puromycin is conferred by the puromycin N-acetyl-transferase gene (pac) from Streptomyces. Puromycin has a fast mode of action, causing rapid cell death at low antibiotic concentrations. Adherent mammalian cells are sensitive to concentrations of 2 to 5 µg /ml, while cells in suspension are sensitive to concentrations as low as 0.5 to 2 µg /ml

### Product Information

<b>Appearance</b>	Liquid
<b>CAS No.</b>	58-58-2
<b>Molecular Formula</b>	$C_{22}H_{29}N_7O_5 \cdot 2HCl$
<b>Chemical Name</b>	3'-[α-Amino-p-methoxyhydrocinnamamido]-3'-deoxy-N,N-dimethyladenosine dihydrochloride
<b>Molecular Weight</b>	156.22
<b>Purity</b>	>98%
<b>Targets</b>	Protein Synthesis

### Storage and Shipping Information

<b>Storage</b>	-80 centigrade
<b>Shipping Conditions</b>	Dry Ice