

Malate dehydrogenase from Bacteria, Recombinant

Cat. No. NATE-1038

Lot. No. (See product label)

Introduction

Description

Malate dehydrogenase is an enzyme in the citric acid cycle that catalyzes the conversion of malate into oxaloacetate (using NAD⁺) and vice versa (this is a reversible reaction). Malate dehydrogenase is not to be confused with malic enzyme, which catalyzes the conversion of malate to pyruvate producing NADPH. Malate dehydrogenase is also involved in gluconeogenesis, the synthesis of glucose from smaller molecules. Pyruvate in the mitochondria is acted upon by pyruvate carboxylase to form oxaloacetate, a citric acid cycle intermediate. In order to get the oxaloacetate out of the mitochondria, malate dehydrogenase reduces it to malate, and it then traverses the inner mitochondrial membrane. Once in the cytosol, the malate is oxidized back to oxaloacetate by cytosolic malate dehydrogenase. Finally, phosphoenol-pyruvate carboxy kinase (PEPCK) converts oxaloacetate to phosphoenol pyruvate.

Synonyms

malic dehydrogenase; L-malate dehydrogenase; NAD-L-malate dehydrogenase; malic acid dehydrogenase; NAD-dependent malic dehydrogenase; NAD-malate dehydrogenase; NAD-malic dehydrogenase; malate NAD dehydrogenase; NAD-dependent malate dehydrogenase; NAD-sp; ECific malate dehydrogenase; NAD-linked malate dehydrogenase; MDH; L-malate-NAD⁺ oxidoreductase; S-malate: NAD⁺ oxidoreductase; EC 1.1.1.37; Malate Dehydrogenase

Product Information

Species	Bacteria
Source	E. coli
Form	Lyophilized powder
EC Number	EC 1.1.1.37
CAS No.	9001-64-3
Molecular Weight	40 kD (SDS-PAGE)
Activity	> 550 units / mg
Contaminants	Fumarase (L-Malate) : < 0.01 % Lactate dehydrogenase : < 0.01 % Aspartate aminotransferase : < 0.01 % Glutamate dehydrogenase (NAD ⁺): < 0.001 % NADH oxidase: < 0.001 %
pH Stability	4.5 - 9.0
Optimum pH	5.5 - 8.0
Thermal stability	<80°C
Optimum temperature	37°C

Storage and Shipping Information

Storage	Below -20°C
----------------	-------------

