

β-Xylosidase from Lactobacillus brevis, Recombinant

Cat. No. NATE-1191

Lot. No. (See product label)

Introduction

DescriptionReleases reducing sugars from birchwood xylan (X0502), also catalyzes the hydrolysis of 4-
methylumbelliferyl-β-D-cellobioside and 4-methylumbelliferyl-β-D-glucopyranoside. This enzyme does
not possess endo-xylanase, arabinoxylanase or β-glucanase activities. β-Xylosidase undergoes post-
translation glycosylation which has been shown to be critical for its proper activity and stability.
Deglycosylation altered the the optimum temperature and pH for activity and decreased its
thermostability.

Synonyms	β-Xylosidase;	β-Xylosidase,	thermostable;	9025-53-0
----------	---------------	---------------	---------------	-----------

Product Information

Source	Lactobacillus brevis ATCC 367
Form	Supplied in 3.2 M ammonium sulphate
EC Number	EC 3.2.1.37
CAS No.	9025-53-0
Molecular Weight	63608.3 Da
Purity	> 95 % as judged by SDS-PAGE
Activity	26.76 U/mg
Concentration	210.83 U/ml
Optimum pH	6
Optimum temperature	40°C
Unit Definition	One unit is defined as the amount of enzyme required to release 1μ mol of oNP per minute from oNP- β -D-xylopyranoside (5 mM) in 20 mM sodium phosphate buffer, pH 6.0, at 35°C, and using an extinction coefficient of 18000 M-1cm-1.

Storage and Shipping Information

Storage Store at 4°C (shipped at room temperature)