

## 3 $\alpha$ -Hydroxysteroid Dehydrogenase from *B. choshinensis*, Recombinant

Cat. No. DIA-413

Lot. No. (See product label)

### Introduction

#### Description

In enzymology, a 3 $\alpha$ -hydroxysteroid dehydrogenase (B-specific) (EC 1.1.1.50) is an enzyme that catalyzes the chemical reaction: androsterone + NAD(P)<sup>+</sup>  $\leftrightarrow$  5 $\alpha$ -androsterane-3,17-dione + NAD(P)H + H<sup>+</sup>. The 3 substrates of this enzyme are androsterone, NAD<sup>+</sup>, and NADP<sup>+</sup>, whereas its 4 products are 5 $\alpha$ -androsterane-3,17-dione, NADH, NADPH, and H<sup>+</sup>. This enzyme belongs to the family of oxidoreductases, specifically those acting on the CH-OH group of donor with NAD<sup>+</sup> or NADP<sup>+</sup> as acceptor, more specifically it is part of the group of hydroxysteroid dehydrogenases.

#### Synonyms

hydroxyprostaglandin dehydrogenase; 3 $\alpha$ -hydroxysteroid oxidoreductase; steroid 3 $\alpha$ ; 3 $\alpha$ -hydroxysteroid dehydrogenase (B-specific); 3 $\alpha$ -hydroxysteroid 3-dehydrogenase (B-specific); 3 $\alpha$ -hydroxysteroid:NAD(P)<sup>+</sup> 3-oxidoreductase (B-specific); EC 1.1.1.50

### Product Information

<b>Species</b>	<i>B. choshinensis</i>
<b>Source</b>	<i>B. choshinensis</i>
<b>Appearance</b>	White lyophilizate
<b>EC Number</b>	EC 1.1.1.50
<b>CAS No.</b>	9028-56-2
<b>Molecular Weight</b>	ca. 41 kDa
<b>Activity</b>	> 30 U/mg lyophilizate
<b>pH Stability</b>	6.0–10.0
<b>Optimum pH</b>	11
<b>Thermal stability</b>	below 45°C
<b>Optimum temperature</b>	50–60°C
<b>Michaelis Constant</b>	2.4 x 10 <sup>-5</sup> M (androsterone) 3.0 x 10 <sup>-6</sup> M (NAD)
<b>Structure</b>	2 subunits of 25 kDa (SDS-PAGE)
<b>Stabilizers</b>	Trehalose
<b>Unit Definition</b>	One unit (U) is defined as the amount of enzyme which produces 1 $\mu$ mol of NADH per min at 25°C and pH 8.9.

### Storage and Shipping Information

<b>Storage</b>	at -20°C
<b>Stability</b>	stable at 37°C for at least four weeks

