

β-Phosphoglucomutase from E. coli, Recombinant

Cat. No. NATE-1251

Lot. No. (See product label)

Introduction

Description

Enzymatically converts β-D-glucose-1-phosphate to β-D-glucose-6-phosphate. β-Phosphoglucomutase enzymatically converts β-D-glucose 1-phosphate to β-D-glucose 6-phosphate. It is involved in starch and sucrose metabolism. This enzyme belongs to the family of isomerases, specifically the phosphotransferases (phosphomutases), which transfer phosphate groups within a molecule. This enzyme participates in starch and sucrose metabolism.

Synonyms

β-phosphoglucomutase; β-D-glucose 1,6-phosphomutase; EC 5.4.2.6

Product Information

Species

E. coli

Source

E. coli

Appearance

White lyophilizate

EC Number

EC 5.4.2.6

CAS No.

68651-99-0

Molecular Weight

ca. 34 kDa

Activity

> 20 U/mg lyophilizate

Contaminants

α-amylase < 5.0 x 10⁻³% α-glucosidase < 5.0 x 10⁻²% NADPH oxidase < 5.0 x 10⁻²%

pH Stability

5.0–9.5

Optimum pH

ca. 7.0

Thermal stability

below 45°C

Optimum temperature

40°C

Michaelis Constant

2.3 x 10⁻⁴ M (β-D-glucose-1-phosphate)

Structure

monomer of ca. 25 kDa (SDS-PAGE)

Activators

Mg²⁺, Mn²⁺, Co²⁺, Ni²⁺

Inhibitors

Hg²⁺, Zn²⁺, Cu²⁺, Cd²⁺

Stabilizers

Lactose, EDTA

Unit Definition

One unit (U) is defined as the amount of enzyme which converts 1 μmol of β-D-glucose-1-phosphate to β-D-glucose-6-phosphate per min at 37°C and pH 7.0.

Storage and Shipping Information

Storage

at -20°C

Stability

Stability (liquid form) stable at 37°C for at least one week Stability (powder form)

Stability

Stability (liquid form) stable at 37 °C for at least one week Stability (powder form)
stable at 30°C for at least one month