

# Immobilized Lipase from *Candida antarctica*

Cat. No. NATE-1260

Lot. No. (See product label)

## Introduction

### Description

Triacylglycerol lipase is an enzyme with system name triacylglycerol acylhydrolase. This enzyme catalyses the following chemical reaction: triacylglycerol + H<sub>2</sub>O ⇌ diacylglycerol + a carboxylate.

### Applications

Lipases are used industrially for the resolution of chiral compounds and the transesterification production of biodiesel.

### Synonyms

EC 3.1.1.3; lipase; triglyceride lipase; tributyrinase; butyryl ester hydrolase; tributyrinase; Tween hydrolase; steapsin; triacetinase; tributyrin esterase; Tweenase; amano N-AP; Takedo 1969-4-9; Meito MY 30; Tweenesterase; GA 56; capalase L; triglyceride hydrolase; triolein hydrolase; tween-hydrolyzing esterase; amano CE; cacordase; triglyceridase; triacylglycerol ester hydrolase; amano P; amano AP; PPL; glycerol-ester hydrolase; GEH; meito Sangyo OF lipase; hepatic lipase; lipazin; post-heparin plasma protamine-resistant lipase; salt-resistant post-heparin lipase; heparin releasable hepatic lipase; amano CES; amano B; tributyrinase; triglyceride lipase; liver lipase; hepatic monoacylglycerol acyltransferase; 9001-62-1

## Product Information

### Source

*Candida antarctica*

### Appearance

Slightly brown

### Form

Beads

### Activity

>2 U/mg

### Unit Definition

1 U corresponds to the amount of enzyme which liberates 1 μmol butyric acid per minute at pH 8.0 and 40°C.

## Storage and Shipping Information

### Storage

2-8°C