

α(2-3) Neuraminidase S from Streptococcus pneumoniae, Recombinant

Cat. No. NATE-1275

Lot. No. (See product label)

Introduction

Description Neuraminidase enzymes are glycoside hydrolase enzymes (EC 3.2.1.18) that cleave the glycosidic linkages of neuraminic acids. Neuraminidase enzymes are a large family, found in a range of organisms. The best-known neuraminidase is the viral neuraminidase, a drug target for the prevention of the spread of influenza infection. The viral neuraminidases are frequently used as antigenic determinants found on the surface of the Influenza virus. Some variants of the influenza neuraminidase confer more virulence to the virus than others. Other homologs are found in mammalian cells, which have a range of functions.

Synonyms neuraminidase; sialidase; α-neuraminidase; acetylneuraminidase; exo-α-sialidase; EC 3.2.1.18; 9001-67-6

Product Information

Species Streptococcus pneumoniae

Source E. coli

Form 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 25°C) and 1 mM EDTA.

Molecular Weight 74000 daltons

Purity > 95% determined by SDS-PAGE

Activity 160,000 units/mg

Unit Definition One unit is defined as the amount of enzyme required to cleave > 95% of the terminal α-Neu5Ac from 1 nmol Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC, in 1 hour at 37°C in a total reaction volume of 10 μl.

Storage and Shipping Information

Storage at -20°C