

## Xylosidase 43A from Bacteroides ovatus, Recombinant

Cat. No. NATE-1527

Lot. No. (See product label)

### Introduction

#### Description

Releases reducing sugars from birchwood xylan (X0502), also catalyzes the hydrolysis of 4-methylumbelliferyl- $\beta$ -D-cellobioside and 4-methylumbelliferyl- $\beta$ -D-glucopyranoside. This enzyme does not possess endo-xylanase, arabinoxylanase or  $\beta$ -glucanase activities.  $\beta$ -Xylosidase undergoes post-translation glycosylation which has been shown to be critical for its proper activity and stability. Deglycosylation altered the the optimum temperature and pH for activity and decreased its thermostability.

#### Synonyms

$\beta$ -Xylosidase; EC 3.2.1.37; 9025-53-0; Xylosidase

### Product Information

#### Species

Bacteroides ovatus

#### Source

E. coli

#### Form

35 mM NaHepes buffer, pH 7.5, 750 mM NaCl, 200 mM imidazol, 3.5 mM CaCl<sub>2</sub>, 0.02% sodium azide and 25% (v/v) glycerol

#### EC Number

EC 3.2.1.37

#### CAS No.

9025-53-0

#### Molecular Weight

39.3 kDa

#### Purity

>90% as judged by SDS-PAGE

#### Concentration

1 mg/mL

#### Optimum pH

6.8

#### Optimum temperature

37 °C

#### Specificity

Xylooligosaccharides

### Storage and Shipping Information

#### Storage

This enzyme is shipped at room temperature but should be stored at -20 °C.