Human glu-Plasminogen

Cat. No. CZY-011

Lot. No. (See product label)

Introduction

Description

Plasminogen is a single chain glycoprotein zymogen which is synthesized in the liver and circulates in plasma at a concentration of approximately 2.4 μM. The plasminogen molecule contains 790 amino acids, 24 disulfide bridges, no free sulfhydryls and 5 regions of internal sequence homology, known as kringles, between Lys77 and Arg560. These five triple-looped, three disulfide bridged, kringle regions are homologous to the kringle domains in t-PA, u-PA and prothrombin. Plasminogen contains one high affinity (Kd=9x10-6M) and four low affinity (Kd=5x10-3M) lysine binding sites. The high affinity binding site resides within the first kringle region of plasminogen. The interaction of plasminogen with fibrin and α2-antiplasmin is mediated by these lysine binding sites. Native glu-plasminogen (Mr=88,000) is readily converted to Lys-77-plasminogen (Mr=83,000) by plasmin hydrolysis of the Lys76-Lys77 peptide bond. Elastase catalyzed cleavage of the Val441-Val442 peptide bond of glu-plasminogen yields a functionally active zymogen termed Val-442 plasminogen or mini-plasminogen. The conversion of plasminogen to plasmin occurs by a variety of mechanisms, but all result in hydrolysis of the Arg560-Val561 peptide bond of plasminogen, yielding two chains which remain covalently associated by a disulfide bond. Native glu-plasminogen is prepared from fresh frozen human plasma by a modification of the procedure of Castellino, utilizing gel filtration and affinity chromatography. The two carbohydrate variants of glu-plasminogen (CHOI and CHOII) are isolated by gradient elution from lysine-Sepharose using the lysine analog, e-aminocaproic acid. The plasminogen is supplied in 50% (vol/vol) glycerol/H2O for storage at -20°C. Purity is determined by SDS-PAGE analysis.

Product Information

Source	Human
Formulation	50% glycerol/water (v/v)
CAS No.	9001-91-6
Molecular Weight	88000
Purity	>95% by SDS-PAGE
Specific Activity	<42 ppm Plasmin activity
Concentration	9.9 mg/mL
Isoelectric point	6.2
Structure	single chain, 24 intra chain disulfide bridges, 5 kringle regions.
Buffer	50% Glycerol/ H2O (vol/vol)
Localization	Plasma
Extinction coefficient	17
Percent carbohydrate	Approximately 2%

Usage and Packaging

Package	1 mg

Storage and Shipping Information

Storage	-20°C
Stability	12 months