

## Human Lys-Plasminogen

Cat. No. CZY-014

Lot. No. (See product label)

### Introduction

#### Description

Plasminogen is a single chain glycoprotein zymogen which is synthesized in the liver and circulates in plasma at a concentration of approximately 2.4  $\mu\text{M}$ . The plasminogen molecule contains 790 amino acids, 24 disulfide bridges, no free sulfhydryls and 5 regions of internal sequence homology, known as kringles, between Lys77 and Arg560. These five triple-looped, three disulfide bridged, kringle regions are homologous to the kringle domains in t-PA, u-PA and prothrombin. Plasminogen contains one high affinity ( $K_d=9 \times 10^{-6}\text{M}$ ) and four low affinity ( $K_d=5 \times 10^{-3}\text{M}$ ) lysine binding sites. The high affinity binding site resides within the first kringle region of plasminogen. The interaction of plasminogen with fibrin and  $\alpha_2$ -antiplasmin is mediated by these lysine binding sites. Native glu-plasminogen ( $M_r=88,000$ ) is readily converted to Lys-77-plasminogen ( $M_r=83,000$ ) by plasmin hydrolysis of the Lys76-Lys77 peptide bond. Elastase catalyzed cleavage of the Val441-Val442 peptide bond of glu-plasminogen yields a functionally active zymogen termed Val-442 plasminogen or mini-plasminogen. The conversion of plasminogen to plasmin occurs by a variety of mechanisms, but all result in hydrolysis of the Arg560-Val561 peptide bond of plasminogen, yielding two chains which remain covalently associated by a disulfide bond. Native glu-plasminogen is prepared from fresh frozen human plasma by a modification of the procedure of Castellino, utilizing gel filtration and affinity chromatography. The two carbohydrate variants of glu-plasminogen (CHOI and CHOII) are isolated by gradient elution from lysine-Sepharose using the lysine analog,  $\epsilon$ -aminocaproic acid. The plasminogen is supplied in 50% (vol/vol) glycerol/H<sub>2</sub>O for storage at -20°C. Purity is determined by SDS-PAGE analysis.

### Product Information

|                               |  |
|-------------------------------|--|
| <b>Source</b>                 | Human  |
| <b>Formulation</b>            | 50% glycerol/water (v/v)   |
| <b>Molecular Weight</b>       | 83000  |
| <b>Purity</b>                 | >95% by SDS-PAGE   |
| <b>Isoelectric point</b>      | 6.7-8.3  |
| <b>Structure</b>              | single chain, 24 intra chain disulfide bridges, 5 kringle regions. |
| <b>Localization</b>           | Plasma   |
| <b>Extinction coefficient</b> | 17   |
| <b>Percent carbohydrate</b>   | Approximately 2%   |

### Usage and Packaging

|                |      |
|----------------|------|
| <b>Package</b> | 1 mg |
|----------------|------|

### Storage and Shipping Information

|                |       |
|----------------|-------|
| <b>Storage</b> | -20°C |
|----------------|-------|

**Storage**

20 °C

**Stability**

12 months