

Native Microorganism Glucose Dehydrogenase (PQQ-dependent)

Cat. No. DIA-192

Lot. No. (See product label)

Introduction

Description

In enzymology, a quinoprotein glucose dehydrogenase (EC 1.1.5.2) is an enzyme that catalyzes the chemical reaction: D-glucose + ubiquinone \leftrightarrow D-glucono-1,5-lactone + ubiquinol. Thus, the two substrates of this enzyme are D-glucose and ubiquinone, whereas its two products are D-glucono-1,5-lactone and ubiquinol. This enzyme belongs to the family of oxidoreductases, specifically those acting on the CH-OH group of donor with a quinone or similar compound as acceptor. This enzyme participates in pentose phosphate pathway. It employs one cofactor, PQQ.

Applications

This enzyme is useful for enzymatic determination of D-Glucose.

Synonyms

Glucose Dehydrogenase; EC 1.1.5.2; D-glucose:ubiquinone oxidoreductase; D-glucose:(pyrroloquinoline-quinone) 1-oxidoreductase; glucose dehydrogenase (PQQ-dependent); glucose dehydrogenase (pyrroloquinoline-quinone); quinoprotein D-glucose dehydrogenase

Product Information

Source

Microorganism

Appearance

Purple amorphous powder, lyophilized

Form

Freeze dried powder

EC Number

EC 1.1.5.2

CAS No.

81669-60-5

Molecular Weight

approx. 100 kDa (by gel filtration)

Activity

Grade III 500 U/mg-solid or more

Contaminants

Glucose dehydrogenase < $1.0 \times 10^{-3}\%$ (NAD-dependent); Hexokinase < $1.0 \times 10^{-3}\%$

pH Stability

pH 3.5-8.5 (25°C, 16hr)

Optimum pH

7

Thermal stability

below 50°C (pH 7.5, 30min)

Optimum temperature

37°C

Michaelis Constant

4.8 mM (D-Glucose)

Inhibitors

Cu⁺⁺, Pb⁺⁺, Ag⁺

Stabilizers

Ca⁺⁺, BSA

Storage and Shipping Information

Stability

Store at -20°C