

Native Microorganism Glutamate Dehydrogenase (NAD-dependent)

Cat. No. DIA-197

Lot. No. (See product label)

Introduction

Description

Glutamate dehydrogenase (GLDH) is an enzyme, present in most microbes and the mitochondria of eukaryotes, as are some of the other enzymes required for urea synthesis, that converts glutamate to α -ketoglutarate, and vice versa. In animals, the produced ammonia is usually used as a substrate in the urea cycle. Typically, the α -ketoglutarate to glutamate reaction does not occur in mammals, as glutamate dehydrogenase equilibrium favours the production of ammonia and α -ketoglutarate.

Applications

This enzyme is useful for enzymatic determination of NH_3 , α -ketoglutaric acid and L-glutamic acid, and for assay of leucine aminopeptidase and urease. This enzyme is also used for enzymatic determination of urea when coupled with urease in clinical analysis.

Synonyms

Glutamate Dehydrogenase; glutamic dehydrogenase; glutamate dehydrogenase (NAD); glutamate oxidoreductase; glutamic acid dehydrogenase; L-glutamate dehydrogenase; NAD-dependent glutamate dehydrogenase; NAD-dependent glutamic dehydrogenase; NAD-glutamate dehydrogenase; NAD-linked glutamate dehydrogenase; NAD-linked glutamic dehydrogenase; NAD-specific glutamic dehydrogenase; NAD-specific glutamate dehydrogenase; NAD: glutamate oxidoreductase; NADH-linked glutamate dehydrogenase; L-glutamate: NAD⁺ oxidoreductase (deaminating); EC 1.4.1.2; GLDH

Product Information

Source

Microorganism

Appearance

White amorphous powder, lyophilized

EC Number

EC 1.4.1.2

CAS No.

9001-46-1

Molecular Weight

approx. 260 kDa

Activity

Gradell 100 U/mg-solid or more

Contaminants

NAD oxidase < $1.0 \times 10^{-2}\%$

Isoelectric point

5.6

pH Stability

pH 5.0-10.0 (25°C, 20hr)

Optimum pH

7.5-8.0 (α -KG \rightarrow L-Glu) 9.0 (L-Glu \rightarrow α -KG)

Thermal stability

below 50°C (pH 8.3, 10min)

Optimum temperature

55°C (α -KG \rightarrow L-Glu) 50°C (L-Glu \rightarrow α -KG)

Michaelis Constant

$9.21 \times 10^{-3}\text{M}$ (NH_3), $4.80 \times 10^{-3}\text{M}$ (α -Ketoglutarate), $7.8 \times 10^{-5}\text{M}$ (L-Glutamate), $1.29 \times 10^{-4}\text{M}$ (NADH), $5.89 \times 10^{-4}\text{M}$ (NAD⁺)

Structure

6 subunits per mol of enzyme

Structure 6 subunits per mol of enzyme

Inhibitors Heavy metals, PCMB, IAA

Storage and Shipping Information

Stability Stable at -20°C for at least one year