

Native *Bacillus* sp. Leucine dehydrogenase

Cat. No. DIA-209

Lot. No. (See product label)

Introduction

Description

In enzymology, a leucine dehydrogenase (EC 1.4.1.9) is an enzyme that catalyzes the chemical reaction: L-leucine + H₂O + NAD⁺ ↔ 4-methyl-2-oxopentanoate + NH₃ + NADH + H⁺. The 3 substrates of this enzyme are L-leucine, H₂O, and NAD⁺, whereas its 4 products are 4-methyl-2-oxopentanoate, NH₃, NADH, and H⁺. This enzyme belongs to the family of oxidoreductases, specifically those acting on the CH-NH₂ group of donors with NAD⁺ or NADP⁺ as acceptor. This enzyme participates in valine, leucine and isoleucine degradation and valine, leucine and isoleucine biosynthesis.

Applications

This enzyme is useful for enzyme determination of L-leucine and the activity of leucine amino-peptidase.

Synonyms

EC 1.4.1.9; Leucine dehydrogenase; L-leucine: NAD⁺ oxidoreductase (deaminating); L-leucine dehydrogenase; L-leucine: NAD⁺ oxidoreductase (deaminating); LeuDH

Product Information

Source

Bacillus sp.

Appearance

White amorphous powder, lyophilized

EC Number

EC 1.4.1.9

CAS No.

9082-71-7

Molecular Weight

245 kDa

Activity

Gradell 20U/mg-solid or more (containing approx. 70% of stabilizers)

Contaminants

Leucylpeptide decomposing enzymes (Leu-Val) < 1.0×10⁻²% (Leu-Gly-Gly) < 1.0×10⁻²% NADH oxidase < 1.0×10⁻²%

pH Stability

pH 5.5-10.5 (25°C, 20hr)

Optimum pH

10.5-10.8 (L-Leu→α-KIC), 9.4 (α-KIC→L-Leu)

Thermal stability

below 60°C (pH 6.9, 10min)

Optimum temperature

above 70°C

Michaelis Constant

1.0×10⁻³M (L-Leucine), 3.9×10⁻⁴M (NAD⁺), 3.5×10⁻⁵M (NADH), 3.1×10⁻⁴M [α-Ketoisocaproate (α-KIC)], 2.0×10⁻¹M (NH₃)

Structure

6 subunits per mol of enzyme

Inhibitors

Na₂S, Hg⁺⁺, Cu⁺⁺, Co⁺⁺, Mg⁺⁺, p-chloromercuribenzoate

Stabilizers

2-Mercaptoethanol, L-cysteine, dithiothreitol, ethylenediaminetetraacetate

Storage and Shipping Information

Stability

Stable at -20°C for at least one year

