

Native Bovine β-Glucuronidase

Cat. No. NATE-0329

Lot. No. (See product label)

Introduction

Description β-glucuronidase catalyzes the breakdown of complex carbohydrates. In humans it converts conjugated

bilirubin into the unconjugated form, making bilirubin suitable for reabsorption.

Applications β -glucuronidase is used as a reporter gene in GUS assays to monitor gene expression.

Synonyms β-glucuronide glucuronohydrolase glucuronidase; exo-β-D-glucuronidase; ketodase; EC 3.2.1.31; 9001-

45-0; β -D-glucuronoside glucuronosohydrolase; GUSB

Product Information

Species Bovine

Source Bovine liver

EC Number EC 3.2.1.31

CAS No. 9001-45-0

Activity Type B-10, \sim 10,000 units/mg solid; Type B-1, > 1,000,000 units/g solid; Type B-3, > 2,000,000 units/g

solid

Pathway Ascorbate biosynthesis, animals, glucose-1P => ascorbate, organism-specific biosystem (from KEGG)

Ascorbate biosynthesis, animals, glucose-1P => ascorbate, conserved biosystem (from KEGG)

Chondroitin sulfate degradation, organism-specific biosystem (from KEGG)

Function AnxA6 is a novel receptor that mediates the endocytosis of bovine beta-glucuronidase.

Unit One unit will liberate 1.0 μg of phenolphthalein from phenolphthalein glucuronide per hr at 37°C at pH

Definition 5.0 (30 min assay).