

## Native Pseudomonas sp. Carboxypeptidase G

Cat. No. NATE-0102

Lot. No. (See product label)

### Introduction

#### Description

Carboxypeptidase G is a lysosomal, thiol-dependent protease, which progressively cleaves  $\gamma$ -glutamyl pteroyl poly- $\gamma$ -glutamate yielding pteroyl- $\alpha$ -glutamate (folic acid) and free glutamate. It is considered highly specific for the  $\gamma$ -glutamyl bond, but not for the C-terminal amino acid of the leaving group.<sup>1</sup> Molecular mass of this homodimer is approximately 90 kDa. The enzyme is activated by  $Zn^{2+}$  ions.

#### Applications

Carboxypeptidase G from *Pseudomonas* sp., or  $\gamma$ -Glutamyl hydrolase, has been used in a study to assess the role of the putidaredoxin COOH-terminus in P-450cam (cytochrome m) hydroxylations. Carboxypeptidase G from *Pseudomonas* sp. has also been used in a study to investigate the effects of nitric oxide on pemetrexed cytotoxicity via NO-cGMP signaling in lung adenocarcinoma cells.

#### Synonyms

$\gamma$ -Glutamyl hydrolase; EC 3.4.17.11; 9074-87-7; glutamate carboxypeptidase; carboxypeptidase G; carboxypeptidase G1; carboxypeptidase G2; glutamyl carboxypeptidase; N-pteroyl-L-glutamate hydrolase

### Product Information

#### Source

*Pseudomonas* sp.

#### Form

lyophilized powder contains sodium acetate salt.

#### EC Number

EC 3.4.17.11

#### CAS No.

9074-87-7

#### Activity

> 3 units/mg protein

#### Composition

Protein, ~70% biuret

#### Unit Definition

One unit will hydrolyze 1.0  $\mu$ mole of L-glutamic acid from (+)amethopterin per min at pH 7.3 at 30°C.

### Storage and Shipping Information

#### Storage

-20°C