

## **Native Porcine Dipeptidyl Peptidase IV**

Cat. No. NATE-0203

Lot. No. (See product label)

## Introduction

Description

Native DPPIV is a ubiquitous type II transmembrane glycoprotein and a serine protease of the S9 prolyloligopeptidase family. In vivo, it is synthesized with a signal peptide, which functions as the membrane anchoring domain. There is an 88% sequence homology between the human and porcine kidney enzymes. Both exist as homodimers with a subunit molecular weight of ~30 kDa. The high mannose 100 kDa DPPIV precursor is processed in the Golgi to yield a 124 kDa heavily N-and O-linked mature glycoprotein. It is then sorted to the apical membrane through the concerted action of both N-and O-linked glycans and its association with lipid microdomains. The porcine enzyme contains 18.3% carbohydrates, which the glycan composition is 0.9% fucose, 3.4% mannose, 5.1% galactose, 8.2% glucosamine, and 0.7% sialic acid. DPPIV is highly expressed on endothelial cells, epithelial cells, and lymphocytes. It is also present in plasma in its soluble form.

Synonyms

EC 3.4.14.5; 54249-88-6; DPPIV; DPP4; dipeptidyl aminopeptidase IV; Xaa-Pro-dipeptidyl-aminopeptidase; Gly-Pro naphthylamidase; postproline dipeptidyl aminopeptidase IV; lymphocyte antigen CD26; glycoprotein GP110; dipeptidyl peptidase IV; glycylproline aminopeptidase; glycylproline aminopeptidase; X-prolyl dipeptidyl aminopeptidase; pep X; leukocyte antigen CD26; glycylprolyl dipeptidylaminopeptidase; dipeptidyl-peptide hydrolase; glycylprolyl aminopeptidase; dipeptidyl-aminopeptidase IV; DPP IV/CD26; amino acyl-prolyl dipeptidyl aminopeptidase; T cell triggering molecule Tp103; X-PDAP

## **Product Information**

**Species** Porcine

**Source** Porcine Kidney

Form Liquid. In 20 mM Tris-HCl, 5 mM CaCl<sub>2</sub>, 1 μM ZnCl<sub>2</sub>, 0.05% NaN<sub>3</sub>, pH 8.0.

**EC Number** EC 3.4.14.5

**CAS No.** 54249-88-6

**Purity** >94% by SDS-PAGE

**Activity** 1.0 U/mg; Specific Activity >40 U/mg protein

 $\textbf{\textit{Unit}} \qquad \qquad \text{One unit is defined as the amount of enzyme that will hydrolyze 1.0 } \mu \text{mole 7-(Gly-Pro)-amino-4-}$ 

**Definition** methylcoumarinamide per min at 37°C, pH 8.5.

## Storage and Shipping Information

**Storage** at -70°C, Avoid freeze/thaw

1/1